arXiv:1512.07594v2 [math.GR] 11 Aug 2016

# FINITE GROUPS WITH 6 OR 7 AUTOMORPHISM ORBITS

ALEX CARRAZEDO DANTAS, MARTINO GARONZI, RAIMUNDO BASTOS

Abstract. Let $G$ be a group. The orbits of the natural action of $\mathrm{Aut}(G)$ on $G$ are called "automorphism orbits" of $G$, and the number of automorphism orbits of $G$ is denoted by $\omega(G)$. In this paper the finite nonsolvable groups $G$ with $\omega(G) \leq 6$ are classified - this solves a problem posed by Markus Stroppel - and it is proved that there are infinitely many finite nonsolvable groups $G$ with $\omega(G) = 7$. Moreover it is proved that for a given number $n$ there are only finitely many finite groups $G$ without nontrivial abelian normal subgroups and such that $\omega(G) \leq n$, generalizing a result of Kohl.

## 1. Introduction

For $G$ a group denote by $\omega(G)$ the number of "automorphism orbits" of $G$, that is, the number of orbits of the natural action of $\mathrm{Aut}(G)$ on $G$. Observe that automorphism orbits are unions of conjugacy classes and hence they give an example of *fusion* in the holomorph $\mathrm{Aut}(G) \ltimes G$, a well-known concept established in the literature (see for example [3, Chapter 7]). It is interesting to ask what can we say about $G$ only knowing $\omega(G)$. It is obvious that $\omega(G) = 1$ if and only if $G = \{1\}$, and it is well-known that if $G$ is a finite group then $\omega(G) = 2$ if and only if $G$ is elementary abelian (this is false in the infinite case: a construction due to Higman, Neumann and Neumann [8, 6.4.6] shows that there is an infinite non-abelian simple group $S$ with $\omega(S) = 2$). Laffey and MacHale in [5] showed that the alternating group $\mathrm{A}_5$ (which has $\omega(\mathrm{A}_5) = 4$) is the only finite nonsolvable group $G$ with $\omega(G) \leq 4$, and Zhang in [13] classified the finite groups in which any two elements of the same order lie in the same automorphism orbit. Markus Stroppel in [10] showed that the only finite nonabelian simple groups $G$ with $\omega(G) \leq 5$ are the groups $\mathrm{PSL}(2, \mathbb{F}_q)$ with $q \in \{4, 7, 8, 9\}$ (see also [11, 2.12] where the classification of finite simple groups is not used), and in [9] and [10] he investigated the properties of the locally compact topological groups $G$ such that $\omega(G)$ (the number of topological automorphism orbits) is finite.

In [9] Stroppel proposed the following problem (Problem 2.5): classify all the finite nonsolvable groups $G$ with $\omega(G) \leq 6$. In this paper we solve Stroppel's problem showing, in Section 2, that if $G$ is a finite nonsolvable group with $\omega(G) \leq 6$ then $G$ is isomorphic to one of $\mathrm{PSL}(2, \mathbb{F}_q)$ with $q \in \{4, 7, 8, 9\}$, $\mathrm{PSL}(3, \mathbb{F}_4)$ or $\mathrm{ASL}(2, \mathbb{F}_4)$. Here $\mathrm{ASL}(2, \mathbb{F}_4)$ is the affine group $\mathrm{SL}(2, \mathbb{F}_4) \ltimes \mathbb{F}_4^2$ where $\mathrm{SL}(2, \mathbb{F}_4)$ acts naturally on $\mathbb{F}_4^2$. Our strategy to prove this result is the following. First, in Lemma 2 we deal with the case in which $G$ is a simple group and in Lemma 3 we deal with the case in which $G$ is an AT-group (that is, any two elements of $G$ of the same order lie in the same automorphism orbit), using a classification theorem proved by Zhang. Then we prove that if $G$ is not simple then it admits a characteristic elementary abelian 2-subgroup $N = {C_2}^n$ (Lemmas 7, 8) and $G/N \cong \mathrm{PSL}(2, \mathbb{F}_4)$

(Lemma 9). In the proof of Lemma 9 we need the notion of CIT group and a nontrivial theorem of Suzuki. Using Lemma 5 we then prove $n \leq 9$, which is a crucial bound that allows us to reduce the problem to a finite and short list of groups, and finally we prove that $N$ must be a minimal normal subgroup of $G$, implying $n = 4$ by Lemma 10. The conclusion follows easily. In our proof we make use of GAP [2] to deal with some small cases.

By a theorem of Landau [6], for a given number $n$ there are only finitely many finite groups with at most $n$ conjugacy classes. The case of automorphism orbits is different, for instance elementary abelian groups all have only two automorphism orbits. However by a result of Kohl [4], for a given number $n$ there are only finitely many finite nonabelian simple groups $S$ with $\omega(S) \leq n$. In Section 3 we generalize this result to the family of all finite groups without nontrivial abelian normal subgroups.

The results presented so far are complemented in Section 4 where we exhibit infinitely many finite nonsolvable groups $G$ with $\omega(G) = 7$. They are the semidirect products (with natural componentwise action) $\mathrm{SL}(2, \mathbb{F}_4) \ltimes V^m$ where $V = \mathbb{F}_4^2$ and $m \geq 2$ (the case $m = 1$ is $\mathrm{ASL}(2, \mathbb{F}_4)$). Actually we prove the more general result

$$\omega(\mathrm{SL}(2, F) \ltimes \mathrm{M}(2 \times m, F)) = 3 + \omega(\mathrm{SL}(2, F))$$

where $\mathrm{M}(2 \times m, F)$ is the $F$-vector space consisting of all the $2 \times m$ matrices with coefficients in $F$, $m \geq 2$, $F$ is any field of characteristic 2 with more than 2 elements and $\mathrm{SL}(2, F)$ acts on $\mathrm{M}(2 \times m, F)$ by multiplication from the left.

In the following, when needed, we make use of the following GAP function that receives as input a group $G$ and computes $\omega(G)$.

```
omega:=function(g);
return Size(OrbitLengthsDomain(AutomorphismGroup(g),g)); end;
```

## 2. Finite nonsolvable groups with $\omega(G) \leq 6$

In this section all considered groups are assumed to be finite. We solve Problem 2.5 of [9]. The following is the main result of this section.

**Theorem 1.** *Let $G$ be a finite nonsolvable group with $\omega(G) \leq 6$. Then $G$ is isomorphic to one of* $\mathrm{PSL}(2, \mathbb{F}_4)$, $\mathrm{PSL}(2, \mathbb{F}_7)$, $\mathrm{PSL}(2, \mathbb{F}_8)$, $\mathrm{PSL}(2, \mathbb{F}_9)$, $\mathrm{PSL}(3, \mathbb{F}_4)$, $\mathrm{ASL}(2, \mathbb{F}_4)$. *Moreover* $\omega(\mathrm{PSL}(2, \mathbb{F}_4)) = 4$, $\omega(\mathrm{PSL}(2, \mathbb{F}_7)) = 5$, $\omega(\mathrm{PSL}(2, \mathbb{F}_8)) = 5$, $\omega(\mathrm{PSL}(2, \mathbb{F}_9)) = 5$, $\omega(\mathrm{PSL}(3, \mathbb{F}_4)) = 6$, $\omega(\mathrm{ASL}(2, \mathbb{F}_4)) = 6$.

For this we need some preparatory results.

A crucial observation, that we will use many times, is that two elements that lie in the same automorphism orbit have the same order; in particular $\omega(G) \geq m$, where $m$ is the number of element orders of $G$, $m = |\{o(g) \ : \ g \in G\}|$.

We list some lemmas.

The following is [9, Lemma 3.2].

**Lemma 1.** *Let $G$ be a group and $N$ a characteristic subgroup of $G$. Then $\omega(G) \geq \omega(N) + \omega(G/N) - 1$.*

The following is a straightforward consequence of Kohl's classification [4, Table 5]. See also [10, Theorem 4.5].

**Lemma 2.** *Let $S$ be a nonabelian simple group with $\omega(S) \leq 6$. Then $S$ is isomorphic to one of $\mathrm{PSL}(2, \mathbb{F}_q)$ for $q \in \{4, 7, 8, 9\}$ or to $\mathrm{PSL}(3, \mathbb{F}_4)$.*

Recall that a group $G$ is called AT-group if whenever $x, y$ are elements of $G$ of the same order, there exists an automorphism $\alpha$ of $G$ such that $x^\alpha = y$. Such groups were classified by Zhang in [13, Theorem 3.1].

**Lemma 3.** *If $G$ is a nonsolvable AT-group and $\omega(G) \leq 6$ then $G$ is isomorphic to one of $\mathrm{PSL}(2, \mathbb{F}_q)$ for $q \in \{4, 7, 8, 9\}$ or to $\mathrm{PSL}(3, \mathbb{F}_4)$.*

*Proof.* If $G$ is simple the result follows from Lemma 2, so now assume $G$ is not simple. We apply Zhang's classification [13, Theorem 3.1]. According to such classification, $G$ is a direct product of groups $M_1 \times \cdots \times M_s$ of pairwise coprime orders, in particular each $M_i$ is characterstic in $G$. Choose the indices so that $M_1$ has even order. Since $G$ is nonsolvable, by the Feit-Thompson theorem $M_1$ is nonsolvable so $\omega(M_1) \geq 4$, hence if $s > 1$ then $\omega(G) \geq \omega(M_1)\omega(M_2) \geq 8$, a contradiction, so $G = M_1$. In cases (1) and (2) of Zhang's theorem $G$ is an extension of solvable groups by the Feit-Thompson theorem (in particular in case (1) the subgroup $H$ is solvable because a cyclic Sylow 2-subgroup yields a characteristic 2-complement), so we are in case (3). Since our $G$ is not simple and $\omega(\mathrm{SL}(2, \mathbb{F}_5)) = 7$, $\omega(\mathrm{SL}(2, \mathbb{F}_7)) = 9$ and $\omega(\mathrm{SL}(2, \mathbb{F}_9)) = 8$ (GAP computation), $G$ is an extension $\mathrm{SL}(2, \mathbb{F}_5) \ltimes B$ where $B$ is abelian of order coprime to $|\mathrm{SL}(2, \mathbb{F}_5)| = 120$, so $B$ is characteristic in $G$ (it is a direct product of Sylow subgroups of $G$) therefore $\omega(G) \geq \omega(G/B) = \omega(\mathrm{SL}(2, \mathbb{F}_5)) = 7$, a contradiction. □

**Lemma 4.** *Assume that $G$ is a nonsolvable group such that $\omega(G) \leq 6$. Then $Z(G) = \{1\}$.*

*Proof.* As $G$ is nonsolvable, by Burnside's $p^a q^b$ theorem (see for example [8, 8.5.3]) there exist three different prime divisors $p_1, p_2, p_3$ of $|G|$. If by contradiction the center $Z = Z(G)$ of $G$ is nontrivial then it contains an element of prime order $p$, and we may assume $p_1 \neq p \neq p_2$. Now $G$ contains elements of orders $1, p, p_1, p_2, p_1p, p_2p$, so since $\omega(G) \leq 6$, the listed orders are precisely all the element orders of $G$ and $G$ is an AT-group. In particular $p = p_3$ and $|G|$ has exactly three prime divisors. If $p$ does not divide $|G/Z|$ then again by Burnside's theorem $G/Z$ is solvable, hence $G$ is solvable, a contradiction. Hence there exists an element $xZ \in G/Z$ of order $p$. Write the order of $x$ in $G$ as $mp^k$ with $(m, p) = 1$. Then $x^m$ has order $p^k$ hence $k = 1$, so that $x^m$ has order $p$. Moreover being $(m, p) = 1$ we have $x^m \notin Z$ being $x \notin Z$. Since $Z$ is characteristic in $G$, there are at least two automorphism orbits of elements of order $p$, a contradiction. □

The following is a key argument to obtain a bound for the order of a nonsolvable group $G$ such that $\omega(G) \leq 6$.

**Lemma 5.** *Let $G$ be a group with a characteristic subgroup $N$ isomorphic to ${C_p}^n$ for some prime $p$. Let $q$ be a prime divisor of $|G/N|$. Assume that*

1. *$N$ contains only one nontrivial automorphism orbit, that is, for all $x, y \in N$ both nontrivial there is $\alpha \in \mathrm{Aut}(G)$ such that $x^\alpha = y$.*
2. *For all $y \in G$ of order $q$ such that $y \notin N$ there is some $x \in N$ such that $xy$ has order $pq$.*
3. *$N \cap Z(G) = \{1\}$.*
4. *The elements of $G$ of order $pq$ are all in the same automorphism orbit.*

*Let $t$ be the number of elements of order $q$ in $G/N$ and suppose $G/N$ can be generated by $\ell$ elements of order $q$. Then $n \leq \ell\lfloor \log_p(t) \rfloor$, where $\lfloor - \rfloor$ denotes the lower integer part.*

*Proof.* Let $S$ be the set of elements of $G/N$ of order $q$. We claim that for each $s \in S$ there exists some $g_s \in G$ of order $pq$ such that $g_s N = s$. Let $s \in S$ and write $s = yN$ with $y \in G$. If $y$ has order $pq$ we choose $g_s = y$. So now assume $y$ does not have order $pq$. Since $N$ is an elementary abelian $p$-group we have $y^{pq} = 1$ hence $y$ has order $p$ or $q$. Observe that $y \notin N$, being $yN$ of order $q$ in $G/N$. If $y$ has order $p$ then $(yN)^p = 1$ so since $yN$ has order $q$ we deduce $p = q$. This proves that in any case $y$ has order $q$. Now we apply hypothesis (2) to find $x \in N$ such that $xy$ has order $pq$. Clearly $xyN = yN$, so we choose $g_s = xy$.

Set $I := \{g_s \ : \ s \in S\}$. We claim that $\bigcup_{i \in I} C_N(i) = N$ and that the $C_N(i)$ are all isomorphic. Let $x \in N$. We look for $j \in I$ such that $x \in C_N(j)$. Let $i \in I$. Since $iN$ has order $q$ in $G/N$, $i^q \in N$. Since $i$ has order $pq$ in $G$, $i^q \neq 1$. Since all nontrivial elements of $N$ lie in the same automorphism orbit there is some $\alpha \in \mathrm{Aut}(G)$ such that $x = (i^q)^\alpha$. Now, clearly $i^q \in C_N(i)$, so $x \in C_N(i)^\alpha = C_N(i^\alpha)$. Observe that being $N$ characteristic, $i^\alpha \notin N$, moreover $(i^\alpha N)^q = (i^q)^\alpha N = N$, so $i^\alpha N$ has order $q$ in $G/N$. Therefore, by the previous point, there is some $j \in I$ such that $i^\alpha N = jN$, in other words $i^\alpha = jy$ for some $y \in N$. We have $x \in C_N(i^\alpha)$, in other words $xjy = jyx$. Now since $xy = yx$ (being $N$ abelian) we deduce $xj = jx$ and this implies $x \in C_N(j)$. The $C_N(i)$ are all isomorphic because if $i, j \in I$ then by hypothesis (4) there is $\alpha \in \mathrm{Aut}(G)$ with $i^\alpha = j$ hence $C_N(j) = C_N(i^\alpha) = C_N(i)^\alpha$.

Now set $p^k := |C_N(i)|$ for all $i \in I$, and $t := |I|$. The fact that $\bigcup_{i \in I} C_N(i) = N$ implies that $t(p^k - 1) + 1 \geq p^n$, and we deduce $p^k \geq (p^n - 1)/t + 1 \geq p^n/t = p^{n - \log_p(t)}$, hence $n - k \leq \lfloor \log_p(t) \rfloor$. Now, let $J$ be a subset of $I$ of cardinality $\ell$ such that $\langle J \rangle N = G$. If $x \in \bigcap_{j \in J} C_N(j)$ then $x$ is central in $\langle J \rangle N = G$ hence $x \in Z(G)$, so $x = 1$ by hypothesis (3). This implies $\bigcap_{j \in J} C_N(j) = \{1\}$ hence

$$p^n = |N| = |N : \bigcap_{j \in J} C_N(j)| \leq \prod_{j \in J} |N : C_N(j)| = p^{\ell(n-k)}$$

which implies $n \leq \ell(n - k) \leq \ell \lfloor \log_p(t) \rfloor$. □

We now prove some technical structural results which we will need in the proof of Theorem 1.

**Lemma 6.** *Let $G$ be a group with a normal subgroup $N$.*

1. *If $xN \in G/N$ has order $p^h$ where $p$ is a prime and $o(x) = mp^k$ with $(m, p) = 1$, then there exists $y \in G$ of order $p^k$ such that $xN = yN$.*
2. *If $N \cong {C_p}^n$ with $n \geq 1$, $p$ an odd prime and $G/N \cong {C_2}^m$ with $m \geq 2$ then $G$ has involutions centralizing elements of order $p$, in particular $G$ has elements of order $2p$.*
3. *If $N$ is an elementary abelian 2-group and $G$ contains an element $a$ inducing an automorphism of order 2 in $N$ then $G$ has elements of order 4 and if $a$ has order 2 then $ay$ has order 4 for some $y \in N$.*

*Proof.* We prove (1). Observe that $x^{p^h} \in N$ and $k \geq h$. If $h = 0$ choose $y = x^m$. Now assume $h > 0$ and write $\alpha m + \beta p^h = 1$ for some $\alpha, \beta \in \mathbb{Z}$. Then $xN = x^{\alpha m + \beta p^h} N = x^{\alpha m} N$ and since $p$ does not divide $\alpha$, $y = x^{\alpha m}$ has order $p^k$.

We prove (2). Let $Q \cong {C_2}^m$ be a Sylow 2-subgroup of $G$ and let $1 \neq a \in Q$. Given $x \in N$, $(xx^a)^a = x^a x = xx^a$ hence $xx^a a$ has order $2p$ if $xx^a \neq 1$. So we may assume $x^a = x^{-1}$ for all $x \in N$ and for all $a \in Q$ different from 1. If $a, b$ are two nontrivial elements of $Q$ then $x^{ab} = (x^{-1})^{-1} = x$ for all $x \in N$, and $ab \neq 1$, so $xab$ has order $2p$ if $x \neq 1$.

We prove (3). We have $a^2 \in C = C_G(N)$. If the order $k$ of $a^2$ is even then a suitable power of $a$ has order 4. So assume that $k$ is odd, and consider $b := a^k$. Then $b^2 = 1$ and, since $k$ is odd, $Ca = Cb$. As $b$ does not centralize $N$, there exists $y \in N$ with $1 \neq byb^{-1}y^{-1} = byby$, and $by$ has order 4. □

The main point of the following two lemmas is to establish the existence of a characteristic elementary abelian 2-subgroup in any nonsolvable non-simple group $G$ with $\omega(G) \leq 6$.

**Lemma 7.** *Assume that the group $G$ is neither solvable nor simple, and that $\omega(G) \leq 6$. Then every minimal characteristic subgroup $M$ of $G$ is elementary abelian.*

*Proof.* Observe that being $M$ a minimal characteristic subgroup of $G$, $M$ is characteristically simple, thus $M \cong S^m$ for some simple group $S$ (see for example [8, 3.3.15]). We need to show that $S$ is abelian, so suppose by contradiction that $S$ is nonabelian. We have $\omega(M) \leq \omega(G) \leq 6$. By [9, 3.17] we have $\omega(M) = \binom{m+\omega(S)-1}{\omega(S)-1}$, so if $m \geq 2$ the condition $\omega(M) \leq 6$ implies $\omega(S) \leq 3$, contradicting [5]. Hence $m = 1$ and $M \cong S$ is simple. By Lemma 2 we then know that $M$ is isomorphic to $S = \mathrm{PSL}(2, \mathbb{F}_q)$ with $q \in \{4, 7, 8, 9\}$. If $C = C_G(M) \neq \{1\}$ then $G$ has a characteristic subgroup isomorphic to $M \times C$ which contains a number of $\mathrm{Aut}(G)$-orbits at least $\omega(M) \cdot \omega(C) \geq 4 \cdot 2$, which is impossible, so $C = \{1\}$ and $G$ is isomorphic to a subgroup $A \leq \mathrm{Aut}(S)$. As $M$ is a proper subgroup of $G$ (being $G$ not simple), we have that $A$ properly contains $S$ (in the guise of inner automorphisms). If $q \neq 9$ then $S$ has prime index in $\mathrm{Aut}(S)$, and $G \cong A = \mathrm{Aut}(S)$ follows. However, we have $\omega(\mathrm{Aut}(S)) \in \{7, 9, 11\}$ in those cases. There remains the case $q = 9$; here $S = \mathrm{PSL}(2, \mathbb{F}_9) \cong \mathrm{A}_6$, and $\mathrm{Aut}(S)/S$ is elementary abelian of order 4. We have $\omega(\mathrm{Aut}(S)) = 13$. The intermediate groups are isomorphic to $\mathrm{PGL}(2, \mathbb{F}_9)$, $\mathrm{S}_6$, and the (non-simple) Mathieu group M(10), with $\omega(\mathrm{PGL}(2, \mathbb{F}_9)) = 8$, $\omega(\mathrm{S}_6) = 8$, $\omega(\mathrm{M}(10)) = 7$. This leads to a contradiction. □

The following table summarizes the nontrivial element orders of $\mathrm{PSL}(2, \mathbb{F}_q)$ for $q \in \{4, 7, 8, 9\}$ for convenience (they will be used in the proofs of lemmas 8 and 9).

| $\mathrm{PSL}(2, \mathbb{F}_4)$ | $\mathrm{PSL}(2, \mathbb{F}_7)$ | $\mathrm{PSL}(2, \mathbb{F}_8)$ | $\mathrm{PSL}(2, \mathbb{F}_9)$ |
|---|---|---|---|
| $2, 3, 5$ | $2, 3, 4, 7$ | $2, 3, 7, 9$ | $2, 3, 4, 5$ |

**Lemma 8.** *Assume that the group $G$ is neither solvable nor simple, and that $\omega(G) \leq 6$. Then every minimal characteristic subgroup $M$ of $G$ is an elementary abelian 2-group.*

*Proof.* Aiming at a contradiction, we consider a minimal counterexample, i.e. a nonsolvable group $G$ with $\omega(G) \leq 6$ and an abelian minimal characteristic subgroup

$N$ which is not an elementary abelian 2-group, with $|G|$ minimal with this property. By Lemma 7, $N \cong {C_p}^n$ for some prime number $p \neq 2$.

(a) $G/N \cong \mathrm{PSL}(2, \mathbb{F}_4)$ and $G$ has elements of order $2p$.

If $G/N$ is not simple then since $\omega(G/N) \leq \omega(G) - 1 \leq 5$, minimality of the counterexample yields a minimal characteristic subgroup $M/N < G/N$ such that $M/N$ is an elementary abelian 2-group, and $\omega(G/M) \leq \omega(G/N) - 1 \leq 4$ yields $G/M \cong \mathrm{PSL}(2, \mathbb{F}_4)$ (by Lemma 2). Observe that $M/N$ is not central in $G/N$ by Lemma 4, hence $|M/N| > 2$, so Lemma 6(2) yields elements of order $2p$ in $M$. Now $M$ contains elements of orders $1, 2, p$ and $2p$, thus $6 \geq \omega(G) \geq \omega(M) + \omega(G/M) - 1 \geq 3 + \omega(\mathrm{PSL}(2, \mathbb{F}_4))$, a contradiction. Therefore $G/N$ is simple, and $\omega(G/N) \leq 5$ implies (by Lemma 2) that $G/N \cong \mathrm{PSL}(2, \mathbb{F}_q)$ with $q \in \{4, 7, 8, 9\}$ so $G/N$ contains an elementary abelian subgroup $H/N$ of order 4, hence Lemma 6(2) yields elements of order $2p$ in $H$. If $G/N \ncong \mathrm{PSL}(2, \mathbb{F}_4)$ then since $G/N$ has four nontrivial prime-power element orders, Lemma 6(1) gives at least seven orbits, contradicting $\omega(G) \leq 6$.

(b) $p \in \{3, 5\}$ and the representatives of the automorphism orbits of $G$ have order 1, $p$ (in $N$) and $2p$, 2, 3, 5 (outside $N$).

The prime divisors of $|G|$ are 2, 3, 5 and $p$, thus $G$ has elements of orders 1, 2, 3, 5, $p$, and $2p$, implying that $p \in \{3, 5\}$ (otherwise $G$ would be an AT-group, contradicting Lemma 3). If the elements of order $p$ were all contained in $N$ then since $p$ divides $|G/N|$ there would be elements of order $p^a$ with $a \geq 2$ inducing elements of order $p$ in $G/N$ (by Lemma 6(1)), thus $G$ would be an AT-group, contradicting Lemma 3.

(c) $n \leq 3\lfloor \log_p(15) \rfloor$.

This is obtained by applying Lemma 5 with $t = 15$ (the number of involutions in $\mathrm{PSL}(2, \mathbb{F}_4)$), $q = 2$ and $\ell = 3$ (being $\mathrm{PSL}(2, \mathbb{F}_4)$ generated by three involutions). Hypotheses (1) and (4) of Lemma 5 hold by point (b) and hypothesis (3) of Lemma 5 holds by Lemma 4. We are left to prove that hypothesis (2) of Lemma 5 holds. Let $y \in G - N$ be an element of order 2 and let $P$ by a Sylow 2-subgroup of $G$, so that $P \cong C_2 \times C_2$ being $|N|$ odd and $G/N \cong \mathrm{PSL}(2, \mathbb{F}_4)$. Applying Lemma 6(2) to the group $NP$ we find an involution $z \in G$ that commutes with some $1 \neq c \in N$. Since the involutions of $G$ are in the same automorphism orbit there is some $\alpha \in \mathrm{Aut}(G)$ such that $y^\alpha = z$. We have $y^\alpha c = c y^\alpha$ and applying $\alpha^{-1}$ we obtain $y c^{\alpha^{-1}} = c^{\alpha^{-1}} y$, so choosing $x = c^{\alpha^{-1}} \in N$, $xy$ has order $2p$.

(d) $p^n \in \{5^2, 3^4\}$.

Let $q = 5$ if $p = 3$ and $q = 3$ if $p = 5$. Let $H$ be a subgroup of $G$ containing $N$ with $|H| = p^n q$. Observe that $G$ does not contain elements of order $pq$ (by point (b)), hence the element orders of $H$ are $1, p, q$. If $k$ is the number of Sylow $q$-subgroups of $H$, we then have $p^n q = |H| = k(q-1) + p^n$ thus $k = p^n$. Sylow's theorem implies $p^n \equiv 1 \mod q$, so if $p = 5$ then $n \leq 3\lfloor \log_5(15) \rfloor = 3$ and $5^n \equiv 1 \mod 3$ implies $n = 2$, and if $p = 3$ then $n \leq 3\lfloor \log_3(15) \rfloor = 6$ and $3^n \equiv 1 \mod 5$ implies $n = 4$.

If $N \cong {C_5}^2$ then we have an embedding of $G/N \cong \mathrm{PSL}(2, \mathbb{F}_4)$ into $\mathrm{GL}(N) = \mathrm{GL}(2, \mathbb{F}_5)$, a contradiction. Now suppose $N \cong {C_3}^4$. Let $K$ be the preimage of a subgroup of $G/N \cong \mathrm{A}_5$ isomorphic to $\mathrm{A}_4$ via the projection $G \to G/N$. Then $K$ has order $81 \cdot 12$. Moreover, if $x \in K - N$ centralizes $N$ then $xN$ belongs to the kernel of the action of $G/N$ on $N$ induced by the conjugation action of $G$, and

since $G/N$ is simple we deduce that $N$ is central in $G$, a contradiction; this implies that $C_K(N) = N$, in particular $N$ equals the Fitting subgroup of $K$. A GAP check shows that there are only two groups of order $81 \cdot 12$ with Fitting subgroup of size 81 and they both have elements of order 9, which is not our case. This leads to a contradiction. □

Recall that a group of even order such that the centralizer of any involution is a 2-group is called CIT group, a class of groups studied by Suzuki in [12]. He proved that if $N$ is the maximal normal 2-subgroup of a CIT group $G$ and $|G/N|$ is even then there are elements of order 2 outside $N$ [12, Theorem 5, page 438]. Observe that if $G$ is not a CIT group then there is an element of odd order $m \neq 1$ centralizing an involution, thus the product of them has order $2m$.

We will use the notion of CIT group in the proof of the following lemma, whose main point is the computation of the quotient of $G$ over the characteristic elementary abelian 2-subgroup established by Lemma 8.

**Lemma 9.** *Let $G$ be a nonsolvable group.*

*(1) If $\omega(G) \leq 5$ then $G \cong \mathrm{PSL}(2, \mathbb{F}_q)$ with $q \in \{4, 7, 8, 9\}$.*
*(2) If $G$ is not simple and $\omega(G) \leq 6$ then $\omega(G) = 6$, there is a characteristic non-central elementary abelian 2-subgroup $N < G$ and $G/N \cong \mathrm{PSL}(2, \mathbb{F}_4)$. Moreover $G - N$ contains elements of order 2 and $G$ contains elements of order 4 .*

*Proof.* Assume that $G$ is neither solvable nor simple, and that $\omega(G) \leq 6$. From Lemma 8 we know that there exists a characteristic non-central elementary abelian 2-subgroup $N < G$.

If $\omega(G) \leq 5$ and $G$ is not simple then $\omega(G/N) = 4$ and $G/N \cong \mathrm{PSL}(2, \mathbb{F}_4)$ contains an involution, which induces by conjugation an automorphism of $N$ of order 2 (because $C_G(N) = N$, being $G/N$ nonabelian simple). Thus Lemma 6(3) secures elements of order 4. Moreover there are elements of order 3 and 5 in $G$ (they are prime divisors of $|G/N|$). So $G$ is an AT-group, contradicting Lemma 3. Thus $G$ is simple and Lemma 2 implies the result. This proves assertion (1).

Now assume $\omega(G) = 6$. Since $\omega(G/N) \leq \omega(G) - 1 = 5$, by assertion (1) we have $G/N \cong \mathrm{PSL}(2, \mathbb{F}_q)$ with $q \in \{4, 7, 8, 9\}$. We split the rest of the proof in items.

(a) If $q \neq 4$ then $G$ contains an element $x$ in $N$ and an element $a \notin N$ with $o(a) = 2$ and $o(ax) = 4$.

This is done by showing that $G$ must be a CIT group. If $G$ is not a CIT group then $G$ has elements of order $2m$ with $m \neq 1$ odd, so since $G/N$ has four nontrivial element orders, all prime powers, and $G$ has elements of order 1 and 2 in $N$, by Lemma 6(1) $G$ has at least seven orbits, a contradiction. This proves that $G$ is a CIT group so by [12, Theorem 5, page 438] $G - N$ contains an element $a$ of order 2 (we can apply this result because since $G/N$ is simple, $N$ is the maximal normal 2-subgroup of $G$). Since $G/N$ is nonabelian simple, $C_G(N) = N$ so $a$ induces an automorphism of $N$ of order 2, implying the existence of elements of order 4 in $G$ of the form $ax$ for some $x \in N$, by Lemma 6(3).

(b) $q = 4$, in other words $G/N \cong \mathrm{PSL}(2, \mathbb{F}_4)$.

Assume by contradiction that $q \neq 4$. $N$ has elements of order 1 and 2, and by the previous point $G$ has elements of order 2 and 4 outside $N$. If $q = 8$ the elements of order 3, 7 and 9 in $G/N$ are induced by prime-power order elements of $G$ by Lemma 6(1), contradicting $\omega(G) = 6$. So $q \neq 8$, in particular $G/N$ has elements of order 4 and $|G|$ has three prime divisors, so there are elements of two distinct odd prime orders, and since $\omega(G) = 6$ this implies that there are representatives of the automorphism orbits of orders 1, 2, 2, 3, 4, $p \in \{5, 7\}$. In particular the elements of order 4 are all in the same automorphism orbit and by Lemma 6(1) the elements of order 4 in $G/N$ are of the form $gN$ with $o(g) = 4$. Using $a, x$ from the previous point we find $g = (ax)^\alpha$ for some $\alpha \in \mathrm{Aut}(G)$. This implies that $gN = (ax)^\alpha N = a^\alpha x^\alpha N = a^\alpha N$ being $N$ characteristic. Therefore $gN$ has order 2 in $G/N$, a contradiction.

(c) $G - N$ contains elements of order 2.

Assume by contradiction that this is not the case. Then $G$ is not a CIT group by [12, Theorem 5, page 438], implying that there are elements of order $2m$ with $m \neq 1$ odd. Moreover, since $G - N$ does not contain elements of order 2, by Lemma 6(1) there is an element $g \in G$ of order $2^b$ with $b \geq 2$ such that $gN$ has order 2 in $G/N$ and the elements of $G/N$ of order 3 and 5 are induced by elements of $G$ of order 3 and 5. Considering the two nontrivial orbits contained in $N$ (represented by elements of orders 1 and 2) we obtain that $G$ is an AT-group, contradicting Lemma 3.

(d) $G$ contains elements of order 4.

Since $G/N$ is nonabelian simple we have $C_G(N) = N$ hence an element of $G - N$ of order 2 (which exists by the previous point) yields an element of $G$ of order 4 by Lemma 6(3).

This concludes the proof. □

**Lemma 10.** *If $n \leq 9$ and $\mathrm{GL}(n, \mathbb{F}_2)$ has an irreducible subgroup isomorphic to $\mathrm{PSL}(2, \mathbb{F}_4)$ then $n = 4$.*

*Proof.* Let $V := {C_2}^n$ and assume $\mathrm{GL}(n, \mathbb{F}_2)$ has an irreducible subgroup $H$ isomorphic to $\mathrm{PSL}(2, \mathbb{F}_4)$. Then the semidirect product $G = H \ltimes V$ is a primitive permutation group of degree $|V| = 2^n$ (see for example [1, Proposition 1.1.12]).

Performing the following with GAP

```
f:=function(n) local l;
l:=AllPrimitiveGroups(DegreeOperation,2^n);
return Filtered(l,x->not(IsSolvable(x)) and Size(x)/(2^n)=60);
end;
List([1..9],f);
```

we find that for $n \leq 9$, $G$ nonsolvable and $|G|/2^n = 60$ imply $n = 4$. □

*Now we proceed to the proof of Theorem 1.* Let $G$ be a nonsolvable group with $\omega(G) \leq 6$. The result follows from Lemma 9 if $\omega(G) \leq 5$, so now assume $\omega(G) = 6$. By Lemma 9 there exists a characteristic elementary abelian 2-subgroup $N$ in $G$ and $G/N \cong \mathrm{PSL}(2, \mathbb{F}_4)$. There are elements of order $1, 2$ in $N$ and elements outside $N$ inducing elements of order $3, 5$ in $G/N$, moreover by Lemma 9 there are elements of order 4 and elements of order 2 outside $N$. This implies that the automorphism orbits of $G$ are represented by elements of orders 1 and 2 in $N$ and 2, 3, 4, 5 outside $N$. This, together with Lemma 6(3) and Lemma 4, implies that we can apply

Lemma 5 with $p = q = 2$, $t = 15$ (the number of involutions in $\mathrm{PSL}(2, \mathbb{F}_4)$) and $\ell = 3$ (being $\mathrm{PSL}(2, \mathbb{F}_4)$ generated by three involutions) finding that $n \leq 3\lfloor \log_2(15) \rfloor = 9$ where $N = {C_2}^n$.

We claim that $N$ is a minimal normal subgroup of $G$. Indeed let $X$ be a minimal normal subgroup of $G$ contained in $N$. Since $N$ has only one nontrivial automorphism orbit, $N$ is generated by the subgroups $X^\alpha$ where $\alpha \in \mathrm{Aut}(G)$, which are minimal normal subgroups of $G$, hence by [8, 3.3.11] applied to $\Omega$ equal to the group of inner automorphisms of $G$, if (by contradiction) $N$ is not a minimal normal subgroup of $G$ then $N$ is a direct product $X \times Y$ where $X, Y$ are normal in $G$ and nontrivial. Any element of $N$ can be written uniquely as $xy$ where $x \in X$ and $y \in Y$. So if $x, y \neq 1$, and $g \in G$ is such that $(xy)^g = xy$ then $x^g y^g = xy$ and by uniqueness of the expression $x^g = x$ and $y^g = y$. This implies that $C_G(xy) = C_G(x) \cap C_G(y)$. Since $C_G(xy) \cong C_G(x) \cong C_G(y)$ (being $x$, $y$, $xy$ in the same automorphism orbit) we obtain $C_G(xy) = C_G(x) = C_G(y)$. This implies that there exists $H \leq G$ such that $C_G(w) = H$ for all $w \in N$ different from 1. Clearly $H$ contains $N$. If $H \neq N$ then there are elements of $G/N \cong \mathrm{PSL}(2, \mathbb{F}_4)$ acting trivially on $N$, a contradiction, so $H = N$. Let $P$ be a Sylow 2-subgroup of $G$: clearly $P$ properly contains $N$. Since $P$ is a finite 2-group and $N \trianglelefteq P$, there is a nontrivial $z \in Z(P) \cap N$, hence $C_G(z) \neq N$, a contradiction.

Being $N$ a minimal normal subgroup of $G$, Lemma 10 implies $n = 4$ and hence $|G| = 2^4 \cdot 60 = 960$. Performing the following with GAP

```
l1:=AllSmallGroups(960);
l2:=Filtered(l1,x-> not(IsSolvable(x)) and Size(Center(x))=1);
List(l2,omega);
```

we find that there is only one nonsolvable group $G$ of size 960 with trivial center and such that $\omega(G) = 6$. This group is $\mathrm{ASL}(2, \mathbb{F}_4)$, as easily verified by computing

```
omega(AllPrimitiveGroups(DegreeOperation,16)[15]).
```

This concludes the proof. □

## 3. Groups without nontrivial abelian normal subgroups

This section gives a generalization of the main result of Kohl's paper [4]. As in that paper, we need the classification of the finite simple groups in our proof.

**Theorem 2.** *Let $n$ be a natural number. Then there are only finitely many finite groups $G$ without nontrivial abelian normal subgroups such that $\omega(G) \leq n$.*

*Proof.* Let $G$ be a finite group without nontrivial abelian normal subgroups, such that $\omega(G) \leq n$, and let $N := \mathrm{soc}(G)$, the socle of $G$, that is, the subgroup of $G$ generated by the minimal normal subgroups of $G$. Suppose that the centralizer $C_G(N)$ (which is a normal subgroup of $G$) is nontrivial and let $M$ be a minimal normal subgroup of $G$ contained in $C_G(N)$. By definition of $N$ we have $M \subseteq N$ hence $M$ centralizes itself, implying that $M$ is abelian, contradicting the fact that $G$ does not have nontrivial abelian normal subgroups. This shows that $C_G(N) = \{1\}$, hence the conjugation action gives an injective homomorphism $G \to \mathrm{Aut}(N)$, implying that $|G| \leq |\mathrm{Aut}(N)| \leq |N|^{|N|}$. This means that it is enough to show that there are only a finite number of possibilities for $N$.

$N$ is a direct product of some minimal normal subgroups by [8, 3.3.11] applied to $\Omega$ equal to the group of inner automorphisms of $G$. Write $N = S_1^{m_1} \times \cdots \times S_t^{m_t}$, where each $S_i$ is a nonabelian simple group and $S_i \ncong S_j$ if $i \neq j$. Observe that each $S_i^{m_i}$ is characteristic in $N$, hence $t \leq \omega(N)$ and $\omega(S_i^{m_i}) \leq \omega(N)$. Moreover $\omega(S_i^{m_i}) = \binom{m_i+\omega(S_i)-1}{\omega(S_i)-1}$ by [9, 3.17], so $m_i \leq \omega(S_i^{m_i})$, $\omega(S_i) \leq \omega(S_i^{m_i})$. Since $\omega(N) \leq \omega(G) \leq n$ being $N$ characteristic in $G$, we deduce $m_i \leq n$, $\omega(S_i) \leq n$ for all $i$ and $t \leq n$. By [4], there are only finitely many possibilities for each $S_i$ hence also for $N$. □

## 4. Infinitely many nonsolvable groups with $\omega(G) = 7$

Let $F$ be a field and $m$ a positive integer. Consider the group $G := G_{m,F} := \mathrm{SL}(2,F) \ltimes V$ where $\mathrm{SL}(2,F)$ (the group of $2\times 2$ matrices of determinant 1 with coefficients in $F$) acts on $V := \mathrm{M}(2\times m, F)$ (the $F$-vector space of $2\times m$ matrices with coefficients in $F$) by multiplication from the left. Observe that $G_{1,\mathbb{F}_4} = \mathrm{ASL}(2,\mathbb{F}_4)$. In the following theorem we compute $\omega(G)$ when $F$ has characteristic 2.

**Theorem 3.** *Let $F$ be a (possibly infinite) field of characteristic 2 with more than 2 elements. If $m \geq 2$ then $\omega(G_{m,F}) = 3 + \omega(\mathrm{SL}(2,F))$.*

*Proof.* We identify $G$ with $\left\{ \begin{pmatrix} X & Y \\ 0 & 1 \end{pmatrix} \ : \ X \in \mathrm{SL}(2,F),\ Y \in \mathrm{M}(2\times m, F) \right\}$. $G$ is a normal subgroup of

$$\Gamma := \left\{ \begin{pmatrix} A & B \\ 0 & C \end{pmatrix} \ : \ A \in \mathrm{GL}(2,F),\ B \in \mathrm{M}(2\times m, F),\ C \in \mathrm{GL}(m,F) \right\}.$$

Observe that $\Gamma$ acts on $G$ by conjugation hence every automorphism orbit of $G$ is a union of $\Gamma$-orbits. In what follows we make use of the identity

$$\begin{pmatrix} A & B \\ 0 & C \end{pmatrix}^{-1} \begin{pmatrix} X & Y \\ 0 & 1 \end{pmatrix} \begin{pmatrix} A & B \\ 0 & C \end{pmatrix} = \begin{pmatrix} A^{-1}XA & A^{-1}(X-1)B + A^{-1}YC \\ 0 & 1 \end{pmatrix}.$$

We identify $V$ with $M = \left\{ \begin{pmatrix} 1 & Y \\ 0 & 1 \end{pmatrix} \ : \ Y \in \mathrm{M}(2\times m, F) \right\}$. Observe that $M$ is the unique maximal abelian normal subgroup of $G$. Indeed, if $K$ is any abelian normal subgroup then $KM/M \cong K/K\cap M$ is an abelian normal subgroup of $G/M \cong \mathrm{SL}(2,F)$ which is simple since $|F| \geq 4$ (note that $\mathrm{SL}(2,F) \cong \mathrm{PSL}(2,F)$ being $F$ of characteristic 2). Thus $KM/M$ is trivial, in other words $K \leq M$. In particular $M$ is characteristic in $G$.

We claim that $M$ contains at least three orbits. Let $v, w \in \mathrm{M}(1\times m, F)$ be linearly independent, let $T_{v,w} = \begin{pmatrix} v \\ w \end{pmatrix}$ and let $g_{v,w} = \begin{pmatrix} 1 & T_{v,w} \\ 0 & 1 \end{pmatrix} \in M$. Observe that $\alpha = \begin{pmatrix} X & Y \\ 0 & 1 \end{pmatrix} \in G$ centralizes $g_{v,w}$ if and only if $(X-1)T_{v,w} = 0$. This implies easily that $C_G(g_{v,w}) = M$. Let $J := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. Both $g_{v,w}$ and $h = \begin{pmatrix} J & T_{v,0} \\ 0 & 1 \end{pmatrix}$ commute with $g_{v,0}$, but $g_{v,w}h \neq hg_{v,w}$. This implies that $C_G(g_{v,0})$ is not abelian. Hence 1, $g_{v,w}$, $g_{v,0}$ have non-isomorphic centralizers, in particular they lie in different automorphism orbits.

We claim that $M$ contains exactly three orbits. By the above, it is enough to prove that $M$ contains at most three orbits. Let $\begin{pmatrix} 1 & Y \\ 0 & 1 \end{pmatrix} \in M$, $e_1 = (1, 0, \ldots, 0)$ and $e_2 = (0, 1, 0, \ldots, 0)$ in $\mathrm{M}(1 \times m, F)$. There exists $C \in \mathrm{GL}(m, F)$ such that $YC = \begin{pmatrix} e_1 \\ 0 \end{pmatrix}$ or $YC = \begin{pmatrix} e_1 \\ e_2 \end{pmatrix}$ according to whether $Y$ has rank 1 or 2. Hence $M$ contains at most three $\Gamma$-orbits, so $M$ contains at most three orbits.

We claim that any non-trivial element of $\mathrm{SL}(2, F)$ of order a power of 2 has order 2, and the elements of order 2 in $\mathrm{SL}(2, F)$ are all conjugated to $J := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ in $\mathrm{GL}(2, F)$. Consider $X \in \mathrm{SL}(2, F)$ of order $2^s$ with $s \geq 1$. Observe that $X^{2^s} = 1$ implies that the eigenvalues $\lambda$ of $X$ verify $\lambda^{2^s} = 1$ so being $F$ of characteristic 2, $(\lambda - 1)^{2^s} = 0$ thus $\lambda = 1$. Since $s \geq 1$, $X$ is conjugated to $J$ in $\mathrm{GL}(2, F)$. In particular $s = 1$.

We claim that $\omega(G) \geq 3 + \omega(G/M)$. $M$ is an elementary abelian 2-group, and $\begin{pmatrix} J & \begin{pmatrix} 0 \\ e_1 \end{pmatrix} \\ 0 & 1 \end{pmatrix}^2 = \begin{pmatrix} 1 & \begin{pmatrix} e_1 \\ 0 \end{pmatrix} \\ 0 & 1 \end{pmatrix}$ has order 2. Thus there are elements of order 4 in $G$, and by the discussion above, in $M$ there are two orbits of elements of order 2. Since $\mathrm{SL}(2, F)$ has elements of order 2 and no elements of order 4, by adapting Lemma 1 we obtain $\omega(G) \geq 3 + \omega(G/M)$.

Each involution $g = \begin{pmatrix} X & Y \\ 0 & 1 \end{pmatrix} \in G - M$ is conjugated to $\begin{pmatrix} J & 0 \\ 0 & 1 \end{pmatrix}$ in $\Gamma$. Indeed, since $X$ is conjugated to $J$ in $\mathrm{GL}(2, F)$, we may assume $X = J$. Then $g^2 = 1$ implies $Y = \begin{pmatrix} v \\ 0 \end{pmatrix}$ with $v \in \mathrm{M}(1 \times m, F)$. Setting $Z := \begin{pmatrix} 0 \\ v \end{pmatrix}$ we obtain $\begin{pmatrix} 1 & Z \\ 0 & 1 \end{pmatrix}^{-1} \begin{pmatrix} J & Y \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & Z \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} J & 0 \\ 0 & 1 \end{pmatrix}$.

We claim that each element $g = \begin{pmatrix} X & Y \\ 0 & 1 \end{pmatrix}$ of order 4 is conjugated to the element $\begin{pmatrix} J & \begin{pmatrix} 0 \\ e_1 \end{pmatrix} \\ 0 & 1 \end{pmatrix}$ in $\Gamma$. Again, we may assume $X = J$, but $Y = \begin{pmatrix} v \\ w \end{pmatrix}$ with $w \neq 0$ now. Using $C \in \mathrm{GL}(m, F)$ with $wC = e_1$ we see that $g$ is conjugated to $\begin{pmatrix} J & Y_1 \\ 0 & 1 \end{pmatrix}$ with $Y_1 = \begin{pmatrix} x \\ e_1 \end{pmatrix}$ for some $x \in \mathrm{M}(1 \times m, F)$. Now $Z := \begin{pmatrix} 0 \\ x \end{pmatrix}$ yields $\begin{pmatrix} 1 & Z \\ 0 & 1 \end{pmatrix}^{-1} \begin{pmatrix} J & Y_1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & Z \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} J & \begin{pmatrix} 0 \\ e_1 \end{pmatrix} \\ 0 & 1 \end{pmatrix}$.

We claim that if $g = \begin{pmatrix} X & Y \\ 0 & 1 \end{pmatrix} \in G$ with $X^2 \neq 1$ (in other words, the order of $g$ is not 1, 2 or 4) then $g$ is conjugated to $\begin{pmatrix} X & 0 \\ 0 & 1 \end{pmatrix}$ in $G$. To prove this observe that

$X^2 \neq 1$ implies that $X-1$ is invertible, because if not then $\ker(X-1)$ is non-trivial, hence $X-1$ is conjugated to a matrix of the form $\begin{pmatrix} 0 & \alpha \\ 0 & \beta \end{pmatrix}$, thus $X$ is conjugated to $\begin{pmatrix} 1 & \alpha \\ 0 & 1+\beta \end{pmatrix}$, and $\det(X) = 1$ implies $\beta = 0$ hence $X^2 = 1$, a contradiction. Set $Z := (1-X)^{-1}Y$. Then $\begin{pmatrix} 1 & Z \\ 0 & 1 \end{pmatrix}^{-1} \begin{pmatrix} X & Y \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & Z \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} X & 0 \\ 0 & 1 \end{pmatrix}$.

Now being $F$ a field of characteristic 2, the automorphism group of $\mathrm{SL}(2,F)$ is generated by the automorphisms induced by conjugation in $\mathrm{GL}(2,F)$ and the field automorphisms (see for example [7, Page 756]). Observe that using conjugation in $\Gamma$ and field automorphism action on each matrix entry of elements of $G$ we see that the elements listed so far form a complete set of representatives of the automorphism orbits of $G$. Therefore $\omega(G) \leq 3 + \omega(\mathrm{SL}(2,F))$ and the result follows. □

Since $\omega(\mathrm{SL}(2,\mathbb{F}_4)) = 4$ we obtain that $\omega(G_{2,\mathbb{F}_4}) = 7$ for all $m \geq 2$. In particular there are infinitely many finite nonsolvable groups $G$ with $\omega(G) = 7$.

## 5. Acknowledgements

We are really grateful to the referee for very carefully reading a previous version of this paper and for the insightful suggestions that contributed greatly to improve and simplify the proofs of Theorems 1, 2, 3. Also, we thank the referee very much for suggesting the current version of Theorem 3.

## References

[1] A. Ballester-Bolinches, L. M. Ezquerro, Classes of Finite Groups; Springer, 2006.
[2] The GAP Group, GAP – Groups, Algorithms, and Programming, Version 4.4; 2005, (http://www.gap-system.org).
[3] D. Gorenstein; Finite Groups. Second edition. Chelsea Publishing Co., New York, 1980.
[4] S. Kohl; Classifying Finite Simple Groups with Respect to the Number of Orbits Under the Action of the Automorphism Group. Comm. Algebra 32 (2004), no. 12, 4785–4794.
[5] T. J. Laffey, D. MacHale; Automorphism Orbits of Finite Groups. J. Austral. Math. Soc. Ser. A 40 (1986), no. 2, 253–260.
[6] E. Landau, Über die Klassenzahl der binaren quadratischen Formen von negativer Diskriminant, Math. Ann. 56 (1903) 671–676.
[7] Loo-Keng Hua, On the automorphisms of the symplectic group over any field. Ann. of Math. (2) 49, (1948). 739–759.
[8] D. J. S. Robinson, A course in the theory of groups. Second edition; Graduate Texts in Mathematics, 80. Springer-Verlag, New York, 1996.
[9] M. Stroppel; Locally compact groups with many automorphisms. J. Group Theory 4 (2001), no. 4, 427–455.
[10] M. Stroppel; Locally compact groups with few orbits under automorphisms. Topology Proc. 26 (2001/02), no. 2, 819–842.
[11] M. Stroppel; Finite simple groups with few orbits under automorphisms. Stuttgart, 2000, http://w5.mathematik.uni-stuttgart.de/fachbereich/Stroppel/Preprints/simple5-preprint.pdf.
[12] M. Suzuki; Finite Groups with Nilpotent Centralizers. Trans. Amer. Math. Soc. 99 1961 425–470.
[13] J. Zhang; On Finite Groups All of Whose Elements of the Same Order Are Conjugate in Their Automorphism Groups. J. Algebra 153 (1992), no. 1, 22–36.